

FCNC in Concurrent Dark Photon & Dark Z Model

Lopamudra Mukherjee

FPCP 2022

University of Mississippi

In Collaboration with Alakabha Datta (UM), Ahmed Hammad (Seoultech), Danny Marfatia (UoH) and Ahmed Rashed (Ain Shams U., Cairo).

May 24, 2022



Introduction

- Hidden dark sector models have been studied in the past to explain low energy anomalies (such as $g - 2$ of muon, Atomic Parity Violation (APV), Higgs decays etc.) [1203.2947](#), [1304.4935](#), [2104.03280](#)
- However these models can also contribute to Flavor Changing neutral currents (FCNC) which have long-standing hints of presence of new physics (NP).
- FCNC decays are sensitive probes of NP.
- Alternately, measurements of FCNC processes can strongly constrain many extensions of the SM such as those of the $U(1)_D$.

FCNC in Dark $U(1)_D$ - Formalism

- We assume that Z_D is associated with a $U(1)_D$ gauge symmetry of a dark sector.
- Z_D gauge boson couples to the SM bosons via kinetic mixing, parametrized by ε , and $Z - Z_D$ mass matrix mixing, parametrized by $\varepsilon_Z = (M_{Z_D}/m_Z)\delta$.

$$\mathcal{L}_{gauge} = -\frac{1}{4}B_{\mu\nu}B^{\mu\nu} + \frac{1}{2}\frac{\varepsilon}{\cos\theta_W}B_{\mu\nu}Z_D^{\mu\nu} - \frac{1}{4}Z_{D\mu\nu}Z_D^{\mu\nu} \quad (1)$$


- As a consequence of the mixing, the Z_D coupling with the SM results into an interaction Lagrangian

$$\mathcal{L}_{int} = \left(-e\varepsilon J_\mu^{em} - \frac{g}{2\cos\theta_W}\frac{M_{Z_D}}{m_Z}\delta' J_\mu^{NC} \right) Z_D^\mu \quad (2)$$


where, $\delta' \simeq \delta + \frac{M_{Z_D}}{m_Z}\varepsilon \tan\theta_W$

FCNC in Dark $U(1)_D$ - Diagrams


$b \rightarrow s Z_D$




1




2




3




4




5




6




7



8



9





10

- The FCNC amplitudes with the dark Z decays were calculated by *F. Xu*, [1504.07415](#) without the dipole term in the amplitude.
- Furthermore, the phenomenology was discussed with the dark boson off shell.
- Here, we fit the parameters of the dark Z model, namely M_{Z_D} , ε & ε_Z , to the available data on the branching fraction of $B \rightarrow K^{(*)}\ell^+\ell^-$ decays in the bins of q^2 .
- We mostly focus in the mass range $0.01 \leq M_{Z_D}/\text{GeV} \leq 2$.

Hadronic Effects in Decay Width

- In the mass range of interest, Z_D can decay into charged or neutral leptons as well as into light hadrons, if kinematically allowed.
- The region $0.5 < M_{Z_D}/\text{GeV} < 2$ is plagued by hadronic resonances such as $\pi\pi$, $\pi^0\gamma$ etc.
- Perturbative QCD fails to provide a reliable way to evaluate vector boson decays into hadrons, so one has to, instead, turn to chiral perturbation theory.
- Earlier analysis did not consider the effect of hadronic resonances in the decay width.
- Using the vector-meson dominance model, one can estimate the decay width to hadrons as done by *Foguel et al., JHEP 04 (2022) 119* for light mediators.



Wilson Coefficients

$$\mathcal{H}_{\text{eff}} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \left[\sum_{i=1\dots 6} C_i \mathcal{O}_i + \sum_{i=7,8,9,10,S,P} (C_i \mathcal{O}_i + C'_i \mathcal{O}'_i) \right] + h.c. \quad (3)$$

$$\mathcal{O}_9 = \frac{e^2}{g^2} (\bar{s} \gamma_\mu P_L b) (\bar{l} \gamma^\mu l), \quad \mathcal{O}'_9 = \frac{e^2}{g^2} (\bar{s} \gamma_\mu P_R b) (\bar{l} \gamma^\mu l) \quad (4)$$

$$\mathcal{O}_{10} = \frac{e^2}{g^2} (\bar{s} \gamma_\mu P_L b) (\bar{l} \gamma^\mu \gamma_5 l), \quad \mathcal{O}'_{10} = \frac{e^2}{g^2} (\bar{s} \gamma_\mu P_R b) (\bar{l} \gamma^\mu \gamma_5 l) \quad (5)$$


$$\begin{aligned} M_{Z_D} &= \langle H_2 | \bar{d}_i \gamma_\mu P_{L/R} d_j | \bar{H}_1 \rangle \left[\{ (E_{c_1, c_2}^{0,A})_{L/R} + (E_{c_1, c_2}^{0,Z})_{L/R} \} g^{\mu\nu} \right. \\ &+ \{ (E_{c_1, c_2}^{2,A})_{L/R} + (E_{c_1, c_2}^{2,Z})_{L/R} \} (g^{\mu\nu} q^2 - q^\mu q^\nu) \left. \right] V_\nu^{Z_D} \\ &+ \langle H_2 | \bar{d}_i i q_\mu \sigma^{\mu\nu} P_{L/R} d_j | \bar{H}_1 \rangle \{ (M_{c_1, c_2}^{1,A})_{L/R} + (M_{c_1, c_2}^{1,Z})_{L/R} \} V_\nu^{Z_D} \quad (6) \end{aligned}$$

Atomic Parity Violation

- A consequence of the sub-GeV dark Z is atomic parity violation.
- The model parameters are tightly constrained from existing data on the weak charge of proton and Caesium atom from parity violating electron scattering experiments.

$$Q_W^{p,SM} = -2g_{AV}^{ep} \left(1 - \frac{\alpha}{2\pi} \right); \quad g_{AV}^{ep} = -1/2 + 2 \sin^2 \theta_W$$


- In presence of the dark Z , $\sin^2 \theta_W \rightarrow \kappa_d \sin^2 \theta_W$ where,
$$\kappa_d = 1 - \varepsilon \left(\delta + \frac{M_{Z_D}}{m_Z} \varepsilon \tan \theta_W \right) \frac{m_Z}{M_{Z_D}} \cot \theta_W f(Q^2)$$



Preliminary Results

Case A : Dark Z model

- The best fit point : $M_{Z_D} = 18.5$ MeV, $\varepsilon = 0.00765$, $\varepsilon_Z = 0.00712$
- Pull wrt SM = 4.7
- $\varepsilon_Z = 0.00712 \Rightarrow \delta \sim 35!$ Problem with APV!




Recall :
 $\varepsilon_Z = (M_{Z_D}/m_Z)\delta$

Preliminary Results

Case B : Add a short distance NP

$$\mathcal{L}_{int} \supset g_{XV} \bar{\mu} \gamma^\mu \mu Z_{D\mu} \quad (7)$$

- This does not affect APV observables.
- For the fit we fix $\delta = 0.1$ and $\varepsilon = 10^{-4}$ and obtain the best fit at : $M_{Z_D} = 15.4$ MeV, $g_{XV} = -0.22$ with a pull of 4.2 from the SM.



Summary & Outlook

- The dark Z model provides a good explanation of the $B \rightarrow K\ell\ell$ data.
- But we need a very large mixing.
- Short distance NP can help overcome this issue.
- We can relax the APV constraints by considering short-distance NP to the first generation fermions.
- We are also exploring the contributions to $B \rightarrow K\nu\nu$ and semileptonic K-meson decays.

Summary & Outlook

- The dark Z model provides a good explanation of the $B \rightarrow K\ell\ell$ data.
- But we need a very large mixing.
- Short distance NP can help overcome this issue.
- We can relax the APV constraints by considering short-distance NP to the first generation fermions.
- We are also exploring the contributions to $B \rightarrow K\nu\nu$ and semileptonic K-meson decays.

THANK YOU!