

20th Conference on Flavor Physics and CP Violation



Hyperon Physics at **BESIII**


Liang Yan

(On behalf of BESIII Collaboration)

Mississippi, 23th-27th May 2022

(online)

Introduction (I)



Hyperons are a laboratory
for strong interaction and
baryon structure

Decay	$\mathcal{B} (10^{-5})$	Events at BESIII
$J/\psi \rightarrow \Lambda\bar{\Lambda}$	189 ± 9	18.9×10^6
$J/\psi \rightarrow \Sigma^+ \bar{\Sigma}^-$	150 ± 24	15.0×10^6
$J/\psi \rightarrow \Xi\bar{\Xi}$	97 ± 8	9.7×10^6
$\psi(2S) \rightarrow \Sigma\bar{\Sigma}$	23.2 ± 1.2	116×10^3
$\psi(2S) \rightarrow \Omega\bar{\Omega}$	5.66 ± 0.30	28×10^3

Introduction (II)

- Polarization
 - Studies of two-body hyperon weak decays plays an important role in the study of the fundamental symmetries P and CP.
 - The polarization of spin $\frac{1}{2}$ hyperon can be determined in two-body weak decays by $(1 + \alpha_0 \mathbf{P}_{\Sigma^+} \hat{\mathbf{p}} / 4\pi)$

Parity violation S state

Parity conservation P state

$$\alpha = \frac{2 \operatorname{Re}(S^* P)}{|S|^2 + |P|^2}, \quad \beta = \frac{2 \operatorname{Im}(S^* P)}{|S|^2 + |P|^2}, \quad \gamma = \frac{|S|^2 - |P|^2}{|S|^2 + |P|^2}$$

α, β, γ could be determined experimentally.

T. D. Lee and C. N. Yang, Phys. Rev. 108, 1645 (1957)


Introduction (II)

- Polarization
 - Studies of two-body hyperon weak decays plays an important role in the study of the fundamental symmetries P and CP.
 - The polarization of spin $\frac{1}{2}$ hyperon can be determined in two-body weak decays by $(1 + \alpha_0 \mathbf{P}_{\Sigma^+} \hat{\mathbf{p}} / 4\pi)$

If the relative phase between hadronic form factor is not zero(Polarized), the decay parameters α_0 and $\bar{\alpha}_0$ could be simultaneous and direct measured, then test CP symmetry.

J/ ψ -> $\Lambda \bar{\Lambda}$

Nature Phys. 15 (2019) 631



Unpolarized $e^+ e^-$ beams -> Transverse polarization:

$$P_y(\cos \theta_\Lambda) = \frac{\sqrt{1 - \alpha_\psi^2} \sin(\Delta\Phi) \cos \theta_\Lambda \sin \theta_\Lambda}{1 + \alpha_\psi \cos^2 \theta_\Lambda}$$

Formulas

$$d\sigma \propto \mathcal{W}(\xi) d\xi \quad \xi = (\theta, \theta_p, \phi_p, \theta_{\bar{p}}, \phi_{\bar{p}})$$

$$\mathcal{W}(\xi) = \mathcal{T}_0(\xi) + \alpha_\psi \mathcal{T}_5(\xi)$$

$$-\alpha_0 \bar{\alpha}_0 \left(\mathcal{T}_1(\xi) + \sqrt{1 - \alpha_\psi^2} \cos(\Delta\Phi) \mathcal{T}_2(\xi) + \alpha_\psi \mathcal{T}_6(\xi) \right)$$

$$+ \sqrt{1 - \alpha_\psi^2} \sin(\Delta\Phi) (\alpha_0 \mathcal{T}_3(\xi) - \bar{\alpha}_0 \mathcal{T}_4(\xi))$$

Phys. Lett. B 772, 16 (2017)

SPIN CORRELATIONS

POLARIZATIONS

$$\mathcal{T}_0(\xi) = 1$$

$$\mathcal{T}_1(\xi) = \sin^2 \theta \sin \theta_p \sin \theta_{\bar{p}} \cos \phi_p \cos \phi_{\bar{p}} + \cos^2 \theta \cos \theta_p \cos \theta_{\bar{p}}$$

$$\mathcal{T}_2(\xi) = \sin \theta \cos \theta (\sin \theta_p \cos \theta_{\bar{p}} \cos \phi_p + \cos \theta_p \sin \theta_{\bar{p}} \cos \phi_{\bar{p}})$$

$$\mathcal{T}_3(\xi) = \sin \theta \cos \theta \sin \theta_p \sin \phi_p$$




$$\mathcal{T}_4(\xi) = \sin \theta \cos \theta \sin \theta_{\bar{p}} \sin \phi_{\bar{p}}$$

$$\mathcal{T}_5(\xi) = \cos^2 \theta$$

$$\mathcal{T}_6(\xi) = \cos \theta_p \cos \theta_{\bar{p}} - \sin^2 \theta \sin \theta_p \sin \theta_{\bar{p}} \sin \phi_p \sin \phi_{\bar{p}}.$$

$J/\psi \rightarrow \Lambda \bar{\Lambda}$

Nature Phys. 15 (2019) 631



$$\langle \alpha \rangle = \frac{\alpha - \bar{\alpha}}{2} = 0.754 \pm 0.003 \pm 0.002$$

CLAS: $\alpha_\Lambda = 0.721 \pm 0.006 \pm 0.005$
PRL 123 (2019) 182301


Parameters	This work	Previous results
α_ψ	$0.461 \pm 0.006 \pm 0.007$	0.469 ± 0.027 BESIII
$\Delta\Phi$ (rad)	$0.740 \pm 0.010 \pm 0.008$	—
α_Λ	$0.750 \pm 0.009 \pm 0.004$	0.642 ± 0.013 PDG
$\bar{\alpha}_\Lambda$	$-0.758 \pm 0.010 \pm 0.007$	-0.71 ± 0.08 PDG

$J/\psi \rightarrow \Lambda \bar{\Lambda}$

arXiv:2204.11058


- 10 Billion J/ψ events are used to update the results.
- The decay parameter a is consistent with previous measurements.
- A_{CP} value is improved with both statistical and systematical uncertainties.

Par.	This work	Previous results [8]
$\alpha_{J/\psi}$	$0.4748 \pm 0.0022 \pm 0.0024$	$0.461 \pm 0.006 \pm 0.007$
$\Delta\Phi$	$0.7521 \pm 0.0042 \pm 0.0080$	$0.740 \pm 0.010 \pm 0.009$
α_-	$0.7519 \pm 0.0036 \pm 0.0019$	$0.750 \pm 0.009 \pm 0.004$
α_+	$-0.7559 \pm 0.0036 \pm 0.0029$	$-0.758 \pm 0.010 \pm 0.007$
A_{CP}	$-0.0025 \pm 0.0046 \pm 0.0011$	$0.006 \pm 0.012 \pm 0.007$
α_{avg}	$0.7542 \pm 0.0010 \pm 0.0020$	-




J/ ψ and $\psi(3686) \rightarrow \Sigma^+ \bar{\Sigma}^-$

Phys. Rev. Lett. 125, 052004 (2020)



$$\frac{dM}{d\cos \theta} \sim \sqrt{1 - \alpha_{\psi}^2} \alpha_0 \sin \Delta\Phi \cos \theta \sin \theta$$

Parameter	Measured value
$\alpha_{J/\psi}$	$-0.508 \pm 0.006 \pm 0.004$
$\Delta\Phi_{J/\psi}$	$-0.270 \pm 0.012 \pm 0.009$
$\alpha_{\psi'}$	$0.682 \pm 0.03 \pm 0.011$
$\Delta\Phi_{\psi'}$	$0.379 \pm 0.07 \pm 0.014$
α_0	$-0.998 \pm 0.037 \pm 0.009$
$\bar{\alpha}_0$	$0.990 \pm 0.037 \pm 0.011$



$$M(\cos \theta) = (m/N) \sum_i^{N_k} (\sin \theta_p^i \cos \phi_p^i - \sin \theta_{\bar{p}}^i \cos \phi_{\bar{p}}^i)$$

The points with error bars are the data, and the solid-line histogram is the global fit result. The dotted histogram is phase space model.

CP asymmetry


$$-0.004 \pm 0.037 \pm 0.010$$

average decay asymmetry

$$-0.994 \pm 0.004 \pm 0.002$$

$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$

arXiv:2105.11155



$$W = \sum_{\mu, \bar{\nu}=0}^3 C_{\mu \bar{\nu}} \sum_{\mu', \bar{\nu}'=0}^3 a_{\mu, \mu'}^{\Xi} a_{\bar{\nu}, \bar{\nu}'}^{\bar{\Xi}} a_{\mu', 0}^{\Lambda} a_{\bar{\nu}', 0}^{\bar{\Lambda}}$$

$d\Gamma \propto W(\xi, \omega)$, ξ : 9 kin. variables
8 parameters:

$$\omega = (\underset{\text{Production}}{\alpha_\Psi, \Delta\Phi, \alpha_\Xi, \phi_\Xi, \alpha_\Lambda, \bar{\alpha}_\Xi, \bar{\phi}_\Xi, \bar{\alpha}_\Lambda}) \underset{\text{Decay}}{}$$

There are 73k events (190 background), the 8 parameters are estimated with unbinned MLL fit!


$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$

arXiv:2105.11155

Parameter	This work	Previous result	
α_ψ	$0.586 \pm 0.012 \pm 0.010$	$0.58 \pm 0.04 \pm 0.08$	[39]
$\Delta\Phi$	$1.213 \pm 0.046 \pm 0.016$ rad	–	
α_Ξ	$-0.376 \pm 0.007 \pm 0.003$	-0.401 ± 0.010	[21]
ϕ_Ξ	$0.011 \pm 0.019 \pm 0.009$ rad	-0.037 ± 0.014 rad	[21]
$\bar{\alpha}_\Xi$	$0.371 \pm 0.007 \pm 0.002$	–	
$\bar{\phi}_\Xi$	$-0.021 \pm 0.019 \pm 0.007$ rad	–	
α_Λ	$0.757 \pm 0.011 \pm 0.008$	$0.750 \pm 0.009 \pm 0.004$	[14]
$\bar{\alpha}_\Lambda$	$-0.763 \pm 0.011 \pm 0.007$	$-0.758 \pm 0.010 \pm 0.007$	[14]
$\xi_P - \xi_S$	$(1.2 \pm 3.4 \pm 0.8) \times 10^{-2}$ rad	–	Independent measurement of α_Λ
$\delta_P - \delta_S$	$(-4.0 \pm 3.3 \pm 1.7) \times 10^{-2}$ rad	$(10.2 \pm 3.9) \times 10^{-2}$ rad [17]	First measurement of weak phase difference!
A_{CP}^Ξ	$(6.0 \pm 13.4 \pm 5.6) \times 10^{-3}$	–	3 CP test
$\Delta\phi_{\text{CP}}^\Xi$	$(-4.8 \pm 13.7 \pm 2.9) \times 10^{-3}$ rad	–	
A_{CP}^Λ	$(-3.7 \pm 11.7 \pm 9.0) \times 10^{-3}$	$(-6 \pm 12 \pm 7) \times 10^{-3}$	[14]
$\langle \phi_\Xi \rangle$	$0.016 \pm 0.014 \pm 0.007$ rad		

$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$

arXiv:2105.11155





Polarization and spin correlations in the $J/\psi \rightarrow \Xi^- \bar{\Xi}^+$


$\psi(3686) \rightarrow \Omega^- \Omega^+$

Phys. Rev. Lett. 126, 092002 (2021)

- The spin of Ω^- $J=3/2$ has never unambiguously confirmed by experiments directly.
- Polarization of the Ω^- can be studied with the Ω^- weak decay chains, and decay parameters could be measured.
- Helicity amplitude method is used.



$\psi(3686) \rightarrow \Omega^- \Omega^+$



Phys. Rev. Lett. 126, 092002 (2021)

3/2 is preferred over 1/2 with significance more than 14σ

Not only observe vector polarization(r_1), but also quadrupole (r_6, r_7, r_8) and octupole(r_{10}, r_{11}) polarizations

$$\begin{aligned} \text{Br}(\psi(3686) \rightarrow \Omega^+ \Omega^-) &= \\ (5.85 \pm 0.12 \pm 0.25) \times 10^{-5} \\ \alpha &= 0.24 \pm 0.10 \end{aligned}$$



Summary

- Hyperons are an important probe to study QCD, fundamental symmetries, and form factors.
- 10 Billion J/ψ data and 2.7 Billion $\psi(3686)$ data collected will bring more exciting results.
- More hyperons study results come soon.

THANK YOU


BACK UP

BEPCII storage rings: a τ -charm factory



Update of BEPC (started 2004, first collisions July 2008)	
Beam energy	1 - 2.47 GeV
Optimum energy	1.89 GeV
Single beam current	0.91 A
Crossing angle	11 mrad
Design luminosity	$1 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
Achieved	$1 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$

BESIII detectors



- Main Drift Chamber (MDC)
 - $\sigma(p)/p = 0.5\%$
 - $\sigma_{dE/dx} = 5.0\%$
- Time-of-flight (TOF)
 - $\sigma(t) = 68\text{ps}$ (barrel)
 - $\sigma(t) = 65\text{ps}$ (endcap)
- Electro Magnetic Calorimeter (EMC)
 - $\sigma(E)/E = 2.5\%$
 - $\sigma_{z,\phi}(E) = 0.5 - 0.7 \text{ cm}$
- RPC MUON Detector
 - $\sigma(xy) < 2 \text{ cm}$